

Alcohol-Induced Deaths by Year, Maine, 1999-2009

Year	Number of Deaths	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
1999	106	8.4	6.8 - 10	7.7	6.2-9.2
2000	97	7.6	6.2 - 9.3	6.9	5.6-8.5
2001	94	7.3	5.9 - 9.0	6.5	5.3-8.0
2002	101	7.8	6.3 - 9.3	7.0	5.6-8.3
2003	117	9.0	7.3 - 10.6	7.7	6.3-9.1
2004	106	8.0	6.5 - 9.6	6.8	5.5-8.1
2005	131	9.9	8.2 - 11.6	8.1	6.7-9.5
2006	114	8.6	7.0 - 10.2	7.4	6.0-8.8
2007	133	10.1	8.4 - 11.8	8.1	6.7-9.5
2008	99	7.5	6.1 - 9.2	6.1	5.0-7.5
2009	133	10.1	8.4 - 11.8	8.4	6.9-9.9

Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death 1999-2009 on CDC WONDER Online Database, released 2012 Program, Maine CDC.

Alcohol-Induced Deaths: Deaths with alcohol-induced causes as the underlying cause of death.

Alcohol-Induced causes: ICD-10 codes: E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, R78.0, X45, X65, and Y15.

Crude rates are deaths per 100,000 population

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population

95% CI: 95% confidence interval of the rate.